




THE LUNG CHANNEL OF HAND TAIYIN



The Lung primary channel



The Lung luo-connecting channel



The Lung divergent channel

LUNG

CHEST	DEADMAN	CAM
Lung 1	On the lateral aspect of the chest, in the first intercostals space, 6 cun lateral to the midline, 1 cun inferior to LU 2	<i>Laterosuperior to the sternum at the lateral side of the first intercostal space, 6 cun lateral to the Ren Meridian</i>
Lung 2	On the antero-lateral aspect of the chest, below he lateral extremity of the clavicle, 6 cun lateral to the midline, in the center of the hollow of the delto-pectoral triangle.	<i>In the depression below the clavicle, 6 cun lateral to the Ren Meridian.</i>
UPPER ARM		
Lung 3	On the antero-lateral aspect of the upper arm, 3 cun inferior to the axillary fold and 6 cun superior to LU 5, in the depression between the lateral border of the biceps brachii muscle and the shaft of the humerus.	<i>On the medial aspect of the upper arm, 3 cun below the end of axillary fold, on the radial side of m. biceps brachii.</i>
Lung 4	On the antero-lateral aspect of the upper arm, 4 cun inferior to the axillary fold and 5 cun superior to LU 5, in the depression between the lateral border of the biceps brachii muscle and the shaft of the humerus.	<i>On the medial aspect of the upper arm, 1 cun below LU 3, on the radial side of m. biceps brachii.</i>

FOREARM

Lung 5	On the cubital crease of the elbow, in the depression at the radial side of the tendon of biceps brachii.	<i>On the cubital crease of the elbow, on the radial side of the tendon of m. biceps brachii. This point is located with elbow slightly flexed.</i>
Lung 6	On the flexor aspect of the forearm, 7 cun proximal to Lu 9, on the line connecting Lu 9 with Lu 5	<i>On the palmer aspect of the forearm, on the line joining Lu 9 and Lu 5, 7 cun above the transverse crease of the wrist.</i>
Lung 7	On the radial aspect of the forearm, approximately 1.5 cun proximal to LI 5, in the cleft between the tendons of brachiorradialis and abductor pollicis longus.	<i>Superior to the styloid process of the radius, 1.5 cun above the transverse crease of the wrist. When the index fingers and thumbs of both hands are crossed with the index finger of one hand placed on the styloid process of the radius of the other, the point is in the depression right under the tip of the index finger.</i>

	DEADMAN	<i>CAM</i>
Lung 8	Above the wrist, 1 cun proximal to Lu 9, on the line connecting Lu 9 with Lu 6, in the depression at the base of the styloid process of the radius and on the radial side of the radial artery.	<i>1 cun above the transverse crease of the wrist in the depression on the lateral side of the radial artery.</i>

WRIST

Lung 9	At the wrist joint, in the depression between the radial artery and the tendon of abductor pollicis longus, level with Ht 7 (the proximal border of the pisiform bone).	<i>At the radial end of the transverse crease of the wrist, in the depression on the lateral side of the radial artery.</i>
---------------	---	---

HAND

Lung 10	On the thenar eminence of the hand, in a depression between the midpoint of the shaft of the first metacarpal bone and the thenar muscles.	<i>On the radial aspect of the midpoint of the first metacarpal bone, on the junction of the red and white skin (ie., the junction of the dorsum and palm of the hand).</i>
Lung 11	On the extensor aspect of the thumb, at the junction of lines drawn along the radial border of the nail and the base of the nail, approximately 0.1 cun from the corner of the nail.	<i>On the radial side of the thumb, about 0.1 cun posterior to the corner of the nail.</i>